



SPORT SCIENCE
INSTITUTE™

INTERASSOCIATION RECOMMENDATIONS

PREVENTING CATASTROPHIC INJURY AND DEATH IN COLLEGIATE ATHLETES



CONTENTS

Introduction 1

Background..... 2

Recommendation 1 | Sportsmanship 5

Recommendation 2 | Protective Equipment 6

Recommendation 3 | Acclimatization and Conditioning 7

Recommendation 4 | Emergency Action Plan 9

Recommendation 5 | Responsibilities of Athletics Personnel..... 10

Recommendation 6 | Education and Training 11

Interassociation Recommendations | Checklist 12

References 16

Appendix A | 2016 Safety in College Football Summit Agenda 18

Appendix B | 2016 Safety in College Football Summit Participants 19

Appendix C | Endorsing Organizations 21



July 2019

NCAA and SSI are trademarks of the National Collegiate Athletic Association.



INTRODUCTION

The second Safety in College Football Summit resulted in interassociation consensus recommendations for three paramount safety issues in collegiate athletics:

1. Independent medical care for collegiate athletes.
2. Diagnosis and management of sport-related concussion.
3. Year-round football practice contact for collegiate athletes.

This document, the fourth arising from the 2016 event, addresses the prevention of catastrophic injury, including traumatic and nontraumatic death, in collegiate athletes. The final recommendations in this document are the result of presentations and discussions on key items that occurred at the summit. After those presentations and discussions, endorsing organization representatives agreed on 18 draft foundational statements (available upon request) that became the basis for this consensus paper, which has been subsequently reviewed by relevant stakeholders and endorsing organizations. This is the final endorsed document for preventing catastrophic injury and death in collegiate athletes.

This document is divided into the following components:

BACKGROUND

This section provides an overview of catastrophic injury and death in collegiate athletes.

INTERASSOCIATION RECOMMENDATIONS: PREVENTING CATASTROPHIC INJURY AND DEATH IN COLLEGIATE ATHLETES

This section provides the final recommendations of the medical organizations for preventing catastrophic injuries in collegiate athletes.

INTERASSOCIATION RECOMMENDATIONS: CHECKLIST

This section provides a checklist for each member school. The checklist will help the athletics health care administrator to ensure that policies are in place and followed, and are consistent with this document, *Interassociation Recommendations: Preventing Catastrophic Injury and Death in Collegiate Athletes*.

REFERENCES

This section provides the relevant references for this document.

APPENDIXES

This section lists the agenda, summit attendees and medical organizations that endorsed this document.

BACKGROUND

Data about catastrophic injuries and illnesses in collegiate athletes began with intermittent accounts from print media, and more formally in 1931, through the American Football Coaches Association's initiation of the Annual Survey of Football Injury Research. Since 1982, the National Center for Catastrophic Sport Injury Research at the University of North Carolina, Chapel Hill,¹ has been the nation's premier source of catastrophic injury and death related to participation in organized sports at all levels of competition, including college. The NCCSIR monitors, collects and analyzes data on catastrophic injuries, illnesses and death and provides publicly available reports about football and other sports.¹

In order to create enhanced national surveillance abilities for catastrophic injuries, illness and death, the NCCSIR has partnered with the Consortium for Catastrophic Injury Monitoring in Sport. The consortium includes the division on traumatic injury at the Matthew Gfeller Sport-Related Traumatic Brain Injury Research Center at the University of North Carolina, Chapel Hill; the division on exertional injury at the Korey Stringer Institute at the University of Connecticut; and the

division on cardiac injury in sport at the University of Washington.¹ Working through the consortium, the NCCSIR has developed new methods of data collection and analysis, including the use of a public-facing online reporting system.²

Researchers who study the epidemiology of catastrophic injury and death in sport identify two mechanisms by which these events occur. *Traumatic* catastrophic injuries, also called direct injuries, are bodily injuries caused directly by participation in a sport activity.¹ An example of a traumatic catastrophic injury is a spinal cord injury caused by tackling in the sport of football. The three leading causes of death from traumatic injury are traumatic brain injuries, spinal cord injuries and internal organ injuries.¹ *Nontraumatic* catastrophic injuries, also known as indirect or exertional injuries, are the "result of exertion while participating in a sport activity or by a complication that was secondary to a non-fatal injury."¹ An example of a nontraumatic catastrophic injury is sudden cardiac arrest in an athlete occurring during a basketball practice. The two leading causes of death from nontraumatic injury are sudden cardiac death and exertional injuries.^{1,3}



Enhancing a culture of safety in college sports in general, and college football in particular, is foundational to reducing the occurrence of catastrophic injury and death and the basis for bringing college athletics stakeholders to the first Safety in College Football Summit in 2014, and then reconvening in 2016. The goal of this and any sport safety initiative is protecting the life and the long-term well-being of all athletes.

Catastrophic Injury Patterns

Since 1982, the first year for which catastrophic injury/illness data were available across all collegiate sports (i.e., NCAA; National Association of Intercollegiate Athletics; National Junior College Athletic Association), there have been 487 catastrophic injuries or illnesses. Of these, 297 (61%) were traumatic events and 190 (39%) were nontraumatic events.¹ In 2016-17, the last year for which data across all collegiate sports are available, 19 catastrophic events occurred, five of which were fatal.¹

Overall, football has the highest number of both traumatic and nontraumatic catastrophic injuries of any collegiate sport. Since 1931, the first year in which football-specific fatality data were collected, there have been 94 traumatic fatalities in college football and 127 nontraumatic fatalities.⁴ More recently, since 1960 there have been 51 traumatic fatalities and 99 nontraumatic in football.⁴ After adjusting for the total number of participating athletes, football is joined by male gymnastics, female skiing, male ice hockey and female gymnastics for the highest rates of traumatic catastrophic injury.¹ Traumatic events in football had fallen every decade from 1960 until 1994. That decline is associated with rule modifications based on research,^{5,6} enhanced medical care and education. Since 1994, the number of traumatic injuries has varied, but at a level generally lower than those of the 1970s and 1980s.¹

Since 1970, in both high school and college football, nontraumatic fatalities have outnumbered traumatic fatalities. Nontraumatic deaths in American football have remained relatively steady for more than five decades. Data from 2017⁴ reveal the current decade will continue this unfortunate and often preventable trend of nontraumatic death that occurs largely in out-of-season or preseason workouts. From 2001 to 2017, the ratio of nontraumatic to traumatic death in collegiate football was 5:1 — 35 nontraumatic deaths compared with seven traumatic fatalities.⁴

While rule modification has the potential to decrease nontraumatic deaths in certain situations (e.g., verification of sickle cell trait decreasing exertional collapse associated with sickle cell trait in Division I football), the policy and procedures to prevent nontraumatic catastrophic death have not kept pace with strength and conditioning sessions and practice sessions that continue to be the setting for record rates of high school and college athlete deaths. For example, of the nine nontraumatic deaths of football players at all levels of the sport in 2017, six occurred during conditioning sessions and one occurred during a strengthening session.¹ For the 2015-16 academic year, six (15%) of the 40 nontraumatic catastrophic injuries and illnesses that occurred across all sports and all levels of competition took place during strength and conditioning sessions.¹ This means that across all sports beside football, nontraumatic injuries are occurring in practice sessions overseen by sport coaches and not during strength and conditioning sessions.

Policy Developments

Available research provides insight into risk factors for catastrophic injury and has led to policy decisions meant to mitigate those risks. Established research demonstrates that NCAA Division I football athletes with sickle cell trait are at a higher risk of nontraumatic catastrophic events, including death.^{5,7,8} In response, the last decade has seen an increase in policy recommendations for the prevention of exertional collapse associated with sickle cell trait (ECAST) in collegiate sport. In 2007, the National Athletic Trainers' Association released a consensus statement on sickle cell trait in the athlete.⁹ By 2013, all three NCAA divisions had adopted legislation requiring confirmation of student-athlete sickle cell trait status before participation.¹⁰⁻¹² This policy, in tandem with targeted on-site precautions, has resulted in a statistically significant decrease in the number of ECAST deaths in college athletes.^{13,14}

Transition periods, defined below, are often associated with poor acclimatization and fitness levels in athletes returning to activity.^{15,16} These concerns have prompted several policy developments. In 2003, the NCAA implemented preseason acclimatization legislation for football.^{17,18} In the same year, NCAA Division I passed a bylaw¹⁹ specific to Football Bowl Subdivision and Football Championship Subdivision football that requires any strength and conditioning

professional who conducts voluntary offseason weight training or conditioning activities to be certified in first aid and cardiopulmonary resuscitation and to be accompanied by a member of the sports medicine staff who has unchallengeable authority to cancel or modify the workout for health and safety reasons. NCAA Division II passed similar legislation one year later.²⁰ In Division I, the unchallengeable authority component of this legislation was extended to all sports other than football in situations when a member of the sports medicine staff is present at a workout. In 2012, NATA released interassociation best practices on the prevention of sudden death in collegiate athletes during strength and conditioning drills.¹⁵ As of 2016, all three NCAA divisions have legislation that requires strength and conditioning professionals to have a certification from either a nationally recognized strength and conditioning certification program^{21,22} or from an accredited strength and conditioning certification program.²³

However, despite these policy developments, catastrophic injuries and fatalities continue to occur. In recent years, most of the fatalities are from nontraumatic causes. These can be mitigated at the member school with appropriate strategies.

Prevention Strategies

Nontraumatic deaths can be mitigated locally through implementation of consensus- and science-based recommendations. Yet, the number of nontraumatic

fatalities are twice those of traumatic fatalities. There have been 99 nontraumatic deaths in collegiate football compared to 51 traumatic deaths since 1960. Just as most of the fatal head injuries and catastrophic cervical spine injuries occurring from 1960 to 1975 can be directly related to the style of play in the sport of football during that time,²⁴⁻²⁶ nontraumatic, exertion-related death is directly related to the conduct and construct of workouts intended to prepare athletes to play sport.^{5,16,27,28} Whereas spearing is often the mechanism for traumatic catastrophic injury and death in football,^{25,26} intense, sustained exertion that is not sport-specific and does not include appropriate work-to-rest ratios, coupled with modifications for individual risk and precautions, is too often the mechanism for exertion-related nontraumatic fatality.^{5,9,16,27-29}

Since 1970, traumatic deaths have undergone a steep and steady decline; nontraumatic deaths, however, have remained steady since 1960. The current era, from 2000 to present, is notable for the following: year-round training for football coupled with the highest incidence of nontraumatic sport-related training deaths in football in recorded history. A proper combination of strategies to prevent the condition from arising in the first place; ensurance of optimal medical care delivery by key stakeholders on-site; and transparency and accountability in workouts should help to eliminate such nontraumatic deaths — a major goal of this document.





INTERASSOCIATION RECOMMENDATIONS:

PREVENTING CATASTROPHIC INJURY AND DEATH IN COLLEGIATE ATHLETES

Best practices for preventing catastrophic injury and death in collegiate sport are organized into six key areas.

RECOMMENDATION 1 | SPORTSMANSHIP

The principle of sportsmanship is foundational to NCAA athletics competition and creates a moral and ethical framework within which athletics competition occurs. This framework rejects any intentional effort by athletes to use any part of their body, uniform or protective equipment as a weapon to injure another athlete or themselves.³⁰ This philosophical commitment is further amplified by expressed statements about the value of sportsmanship in sport playing rules.³¹⁻³³ The avoidance of on-field or on-court behaviors intended to cause injury to another athlete must become part of the cultural foundation from which all subsequent sport safety initiatives arise.

While acknowledging that football, like other contact/collision sports, is an aggressive, rugged contact sport, the rules of football and of all other sports identify a responsibility shared by all involved to conduct themselves according to a shared ethical code.³³ This code requires that the head and helmet not be used as a weapon, and that unsportsmanlike efforts to deliberately injure an opponent are outside the boundaries of fair and legal play. The act does

not need to be purposeful to be considered an infraction.

Given this commitment to sportsmanship, coupled with the considerable safety implications of its violation, the following recommendations regarding deliberate injury to an opponent should be considered in all sports:

1. A player should be ejected immediately from competition (in addition to a particular penalty) for a first infraction.
2. Video replay (when available) after the competition can verify missed calls and could lead to suspension from the next competition. Conferences play a crucial role in this process and should commit themselves to this responsibility.
3. Officials who fail to call such infractions should be educated and/or disciplined appropriately.
4. In helmeted sports, rules should be further developed to prohibit and penalize the initiation of contact with the head/helmet and should be uniformly enforced.



RECOMMENDATION 2 | PROTECTIVE EQUIPMENT

Protective equipment that is used in sport typically must be manufactured and maintained according to performance and safety standards promulgated by standards organizations such as the National Operating Committee on Standards for Athletic Equipment^{34,35} and ASTM International.³⁶ When sport playing rules require equipment to comply with existing standards, the legality of the equipment is dependent on compliance, certification or both with existing standards. For example, current playing rules in the sport of football require that helmets be manufactured and maintained according to standards established by NOCSAE. These standards have been demonstrated to reduce the occurrence of

catastrophic brain injury.³⁷ In some cases, as with the helmet in the sport of football, equipment must be maintained through a reconditioning process. Where this responsibility exists, member institutions must remain vigilant about ensuring necessary maintenance to ensure the continued safety and legality of protective equipment.

The following should be implemented across all sports:

- Every member school should establish policy to ensure annual certification, recertification and compliance, as appropriate, with all protective equipment standards.

RECOMMENDATION 3 | ACCLIMATIZATION AND CONDITIONING

Many nontraumatic deaths take place during the first week of activity of a transition period in training.¹⁵ Given this fact, it is imperative to recognize the vulnerability during these periods and to ensure that both proper exercise and heat acclimatization are implemented.

Transition periods hold particular risk, but absent adherence to established standards, best practices and precautions, collegiate athletes are at risk at all points in the offseason regimen. For example, February and July typically are not transition times, yet from 2000 to 2017, they are the deadliest months of winter and summer training in collegiate football.²⁸ Acclimatization and physiologic progression with a basis of exercise science and sport specificity are the cornerstones of safe conditioning and physical activity. It takes approximately seven to 10 days for the body to acclimatize to the physiologic and environmental stresses placed upon it at the start of a conditioning or practice period, especially during periods of warm or hot weather.^{15,38,39}

Acclimatization, especially heat acclimatization, can occur only through repeated exposure to a hot environment⁴⁰ while progressively increasing the volume and intensity of physical activity.⁴¹ Unfortunately, perceived time pressures by coaches coupled with the culture of certain sports that excesses in training make athletes tough, disciplined and accountable contribute to a tendency to overload athletes during transition periods.^{15,42}

A minimum expectation is that *all* strength and conditioning sessions, regardless of when in the year they occur, should be evidence- or consensus-based; sport-specific; intentionally administered; appropriately monitored, regardless of the phase of training; and not punitive in nature.

For acclimatization and conditioning, the following direction should be considered for all sports and by any individual responsible for the planning and/or implementation of training and conditioning sessions, whether that be a strength and conditioning professional or a sport coach:

1. Training and conditioning sessions should be introduced intentionally, gradually and progressively to encourage proper exercise

acclimatization and to minimize the risk of adverse effects on health. This is especially important during the first seven days of any new conditioning cycle, which should be considered a *transition period*. A lack of progression and sport-specificity in the volume, intensity, mode and duration of conditioning programs in transition periods has been noted as a primary factor in nontraumatic fatalities.¹⁶ Importantly, in this period of year-round sport, new conditioning cycles can occur several times throughout the year and are not limited to the beginning of a competitive season. During transition periods, athletes should be instructed to avoid additional volunteer sessions of physical activity (e.g., 7-on-7 drills, pickup games, drill work). Physical activity schedules during transition periods should be well prescribed, accounting for all sources of physical activity in which an athlete engages.

Examples of transition periods for athletes include, but are not limited to:

- a. Individual transitions.
 - (1) Athletes new to the program.
 - (2) Returning after an injury or illness.
 - (3) Any delayed participation relative to the team schedule.
 - b. Team transitions.
 - Resumption of training after an academic break (e.g., winter, spring, summer breaks).
2. Training and conditioning sessions should be exercise-science based and physiologically representative of the sport and its performance components. Conditioning programs should begin with work-to-rest ratio intervals appropriate for the goals of the training session and that allow for proper recovery.
 3. Collegiate athletes are especially vulnerable to exertional injuries during the first four days of transition periods, and the data support that modifications in these periods can greatly decrease the risk of catastrophic events.¹⁴ During this time, training and conditioning sessions should be appropriately calibrated and include

limitations on total volume and intensity of activity. This may be accomplished in several ways. For example, holding only one training and conditioning session per day during the transition period may be effective for limiting the volume of physical activity.

- a. Properly training during transition periods also should greatly reduce or eliminate rhabdomyolysis, which is largely preventable. Since 2007, 57 NCAA collegiate athletes have been reported as suffering from exertional rhabdomyolysis in nine team outbreaks representing eight different institutions, with 51 of the afflicted athletes requiring hospitalization.⁴³ Novel overexertion, or exertion caused by new activities or at unaccustomed volume or intensity, is the most common cause of exertional rhabdomyolysis and is characterized as too much, too soon and too fast in a workout regimen.¹⁴ Team outbreaks of exertional rhabdomyolysis in NCAA athletes have similarities of irrationally intense workouts designed and conducted by coaches and/or strength and conditioning professionals.⁴³
- b. When phasing in activity during transition periods, athletics staff members should consider the following:
 - (1) Days/week.
 - (2) Body part.
 - (3) Activity/exercise.
 - (4) Sets/repetitions/distance.
 - (5) Load (percent of one-repetition maximum, i.e., 1RM).
 - (6) Work-rest ratio.
 - (7) Modifications: position; individual; return from injury; environment.

4. All training and conditioning sessions should be documented. In addition, all training and conditioning sessions should:
 - a. Be approved by a credentialed strength and conditioning professional, or by the head sport coach at institutions that do not employ strength and conditioning professionals.
 - b. Address exercise volume, intensity, mode and duration.
 - c. Ensure the location of the training and conditioning session is identified in the plan to accommodate venue-specific emergency action planning.
 - d. Be reproducible upon request and be shared with the primary athletics health care providers (team physician and athletic trainer) before the session in which they are to be used.
 - e. Be modified in response to hazardous environmental conditions, scheduling considerations, etc. The amended workout plan should maintain the above principles.
5. A disciplinary system should be developed and applied to strength and conditioning professionals and sport coaches who fail to follow these recommendations. Such penalties could include suspension and/or termination by the member school. Additionally, failure to follow the recommendations could be a reportable offense by member schools to the NCAA.

RECOMMENDATION 4 | EMERGENCY ACTION PLAN

There is broad agreement that the most effective way to prevent catastrophic fatalities and manage nonfatal catastrophic events is through a sound and well-rehearsed emergency action plan.^{13,15,29,44,45} Venue-specific emergency action plans are a cornerstone of emergency readiness for campus and athletics health care providers.^{38,46}

Emergency action plans should be readily available to all members of the athletics community, located both centrally and at each venue at which athletics activities will occur and should be rehearsed with all relevant sports medicine and coaching staff at least once a year. The equipment necessary to execute the emergency action plan should be available to each venue at which athletics activities will occur. Emergency action plan rehearsal also should be incorporated into new employee orientation.

At a minimum, well-rehearsed and venue-specific emergency action plans should be developed for the following nontraumatic catastrophic events:

1. Head and neck injury.
2. Cardiac arrest.
3. Heat illness and heat stroke.
4. Exertional rhabdomyolysis.
5. Exertional collapse associated with sickle cell trait.
6. Any exertional or nonexertional collapse.
7. Asthma.
8. Diabetic emergency.
9. Mental health emergency.

In addition, well-rehearsed and venue-specific emergency action plans should be consistent with the NCAA Concussion Safety Protocol Checklist.⁴⁷ This checklist was created in response to NCAA legislation passed by the Division I conferences with autonomy in January 2015,⁴⁷ and subsequently by all three divisions. The checklist facilitates the development of a comprehensive and coordinated set of policies to guide institutions in the diagnosis and management of collegiate athlete concussions and in the eventual return to play and return to the classroom by those athletes. Concussion emergency action plans should be created for the following suspected conditions:

1. Concussion.
2. Moderate or severe traumatic brain injury.

3. Cervical spine injuries.

Special considerations:

1. Cardiac emergencies — Research has shown that in sudden cardiac arrest, the probability of survival drops by 7-10% for every minute of active arrest, whereas the probability of survival is 89% in properly administered CPR and automated external defibrillators.^{48,49} The location of AEDs should be documented and should reflect a strategy that ensures their arrival at the scene of a collapse with the target goal of collapse-to-shock in less than three minutes.⁴⁹ All AEDs should be checked at least monthly to assure they are fully charged.⁴⁹
2. Exertional heat illness emergencies — Exertional heatstroke is a medical emergency that is characterized by extreme hyperthermia (>40.0 degrees C/>104 degrees F) and central nervous system dysfunction such as altered behavior or decreased consciousness.⁴¹ To differentiate heatstroke from other acute medical events, primary athletics health care providers should be prepared to measure core body temperature using rectal thermometry. Rectal temperature has been demonstrated as the most accurate method for measuring body temperature, whereas other methods such as axillary, tympanic (aural), temporal, oral and skin measurements are not valid or reliable predictors of core temperature.⁵⁰ During warm weather events, but especially pre-season practices of fall season sports, resources (e.g., equipment and personnel) should be readily available to ensure that full-body ice water immersion can be conducted in a timely manner. Full-body immersion in cold water (1.7 degrees C to 15.0 degrees C/35 degrees F to 59 degrees F) is the most effective immediate treatment of exertional heatstroke, with fatality rates close to zero if the body temperature is brought to less than 40.0 degrees C within 30 minutes after collapse.⁴¹ Full-body cold water immersion should be conducted before patient transport, and should be continued until the body has cooled to a temperature below 38.9 degrees C/102 degrees F. During cold water immersion, body temperature should be continuously monitored with rectal thermometry.

RECOMMENDATION 5: RESPONSIBILITIES OF ATHLETICS PERSONNEL

Physical activity never should be used for punitive purposes. Exercise as punishment invariably abandons sound physiologic principles and elevates risk above any reasonable performance reward.¹⁵ As stated in the 2014-15 NCAA Sports Medicine Handbook, this principle has been reinforced by the NCAA Committee on Competitive Safeguards and Medical Aspects of Sports.⁵¹ All athletics personnel, including both sport and strength and conditioning professionals, as well as primary athletics health care providers, should intervene when they suspect that physical activity is being used as punishment. Although “intent” of punishment may be difficult to establish, punishment workouts use unsound physiological principles, as enumerated in this document.

All training and conditioning sessions should be administered by personnel with demonstrated competency in the safe and effective development and implementation of training and conditioning activities, and with the necessary training to respond to emergency situations arising from those activities.

NCAA bylaws in all three divisions require that strength and conditioning professionals have a strength and conditioning certification from either a nationally accredited²³ or nationally recognized,^{21,22} strength and conditioning certification program. Additional NCAA bylaws in Division I¹⁹ require that strength and conditioning professionals must be accompanied by members of the sports medicine staff when conducting voluntary, offseason conditioning sessions. In these situations, NCAA bylaws in both Divisions I and II^{19,20} require the sports medicine staff members have unchallengeable authority to cancel or modify workouts for health and safety reasons.

In Division III, where the presence of full-time strength and conditioning professionals may be less frequent, and where as a result, sport coaches may provide strength and conditioning services to all collegiate athletes, legislation is more nuanced. Any sport coach can conduct an in-season workout without needing a strength and conditioning certification. Only strength and conditioning professionals with nationally recog-

nized certifications can conduct voluntary workouts in the offseason, and then only during the regular academic year and only if the voluntary workouts are being conducted for all collegiate athletes.²¹ This legislation anticipates a situation when a sport coach is otherwise serving a broader, campus-wide responsibility as strength and conditioning beyond the sport he or she coaches.

The following questions about the strength and conditioning credential should be considered when hiring a strength and conditioning professional:

1. Is the strength and conditioning credential one that reflects attaining relevant competencies in the delivery of strength and conditioning services to collegiate athletes and teams?
2. Is the credential conferred by a certification program/process that is nationally accredited?
3. What are the requisite educational standards required for certification eligibility, and the continuing education requirements required by the certification program?
4. Does the certification require CPR and AED certification?
5. Does the certification require a baccalaureate degree or higher, and is it in a degree field with relevance to the provision of strength and conditioning services?

The current state of credentialing across the strength and conditioning profession makes it difficult to ensure that all strength and conditioning professionals have the requisite competency to safely and effectively conduct conditioning sessions. Many organizations currently offer “strength and conditioning” credentials, though there is significant variability in both the content represented by these credentials and the rigor required to attain them. The complete absence of state regulation further complicates this landscape because there is no clearly established strength and conditioning scope of practice, and therefore, there is no authoritative accounting of the knowledge and skill domains required for the safe and effective practice of a strength and conditioning professional. If carefully considered, the five questions above can assist

institutions in identifying strength and conditioning credentials reflecting the attainment of minimal competence in the provision of strength and conditioning services. Moreover, the U.S. Registry of Exercise Professionals (see usreps.org/Pages/Default.aspx) contains those strength and conditioning professions with certifications from programs accredited by the National Commission for Certifying Agencies. NCCA accreditation is considered a marker of quality for certification programs in the health and/or medical domains.

An additional problem arises through the increasingly close alignment between sport coaches and strength and conditioning professionals, especially in the sport

of football. Strength and conditioning professionals frequently are hired by the head football coach, and/or subject to their administrative oversight. This alignment is problematic because it contributes to the perception that strength and conditioning professionals are members of the coaching staff rather than independently credentialed strength and conditioning professionals. Such singular alignment and reporting are not consistent with this document. All strength and conditioning professionals should have a reporting line into the sports medicine or sport performance lines of the institution. This includes sport coaches who have responsibility for providing strength and conditioning services across all sport teams.

RECOMMENDATION 6 | EDUCATION AND TRAINING

Beyond strength and conditioning professionals, each institution should adopt requirements for the education and training of athletics personnel, including as a minimum, but not limited to, strength and conditioning professionals, sport coaches and primary athletics health care providers. Education should focus on preventing catastrophic injury and sudden death in sport. Such education and training should occur annually. Regular education not only can serve to improve the recognition and response skills of those who may be involved in a catastrophic event but also can contribute to a heightened state of organizational mindfulness that contributes to an environment of emergency readiness. Education and prevention strategies should be customized for the unique learning needs of relevant stakeholders and their roles on the athletics team.

Such training should include the following:

1. Foundational information regarding emergency action plans.
2. Environmental monitoring (heat/humidity, lightning).
3. Head and neck injuries.
4. Cardiac arrest.
5. Heat illness and heatstroke.
6. Exertional Rhabdomyolysis.
7. Exertional collapse associated with sickle cell trait.
8. Any exertional or nonexertional collapse.
9. Asthma.
10. Diabetic emergency.
11. Mental health emergency.
12. Proper training principles/principles of periodization.





SPORT SCIENCE
INSTITUTE™

INTERASSOCIATION RECOMMENDATIONS | CHECKLIST

PREVENTING CATASTROPHIC INJURY AND DEATH IN COLLEGIATE ATHLETES

This checklist will help the athletics health care administrator to ensure that policies are in place and followed, and are consistent with this document, *Interassociation Recommendations: Preventing Catastrophic Injury and Death in Collegiate Athletes*.

1 TRAUMATIC: GENERAL	YES	NO	COMMENTS
In all sports, all practices and competitions adhere to existing ethical standards.	<input type="checkbox"/>	<input type="checkbox"/>	
In all sports, using playing or protective equipment as a weapon is prohibited during all practices and competitions.	<input type="checkbox"/>	<input type="checkbox"/>	
In all practices and competitions, deliberately inflicting injury on another player is prohibited.	<input type="checkbox"/>	<input type="checkbox"/>	
All playing and protective equipment, as applicable, meets relevant equipment safety standards and related certification requirements.	<input type="checkbox"/>	<input type="checkbox"/>	
There is a regularly rehearsed emergency action plan consistent with the Concussion Safety Protocol Checklist for all venues at which practices or competitions are conducted.	<input type="checkbox"/>	<input type="checkbox"/>	
There is a regularly rehearsed emergency action plan consistent with the Concussion Safety Protocol Checklist for all suspected concussions.	<input type="checkbox"/>	<input type="checkbox"/>	
There is a regularly rehearsed emergency action plan consistent with the Concussion Safety Protocol Checklist for all suspected moderate or severe traumatic brain injuries.	<input type="checkbox"/>	<input type="checkbox"/>	
There is a regularly rehearsed emergency action plan consistent with the Concussion Safety Protocol Checklist for all suspected cervical spine injuries.	<input type="checkbox"/>	<input type="checkbox"/>	
Annual education and prevention strategies about catastrophic injuries are provided to all sports coaches.	<input type="checkbox"/>	<input type="checkbox"/>	
Annual education and prevention strategies about catastrophic injuries are provided to all strength and conditioning professionals.	<input type="checkbox"/>	<input type="checkbox"/>	

TRAUMATIC: GENERAL CONTINUED	YES	NO	COMMENTS
Annual education and prevention strategies about catastrophic injuries are provided to all primary athletics health care providers (i.e., team physicians and athletic trainers).	<input type="checkbox"/>	<input type="checkbox"/>	
Annual education and prevention strategies about catastrophic injuries are provided to all collegiate athletes.	<input type="checkbox"/>	<input type="checkbox"/>	
Annual education and prevention strategies about catastrophic injuries are provided to all athletics administrators.	<input type="checkbox"/>	<input type="checkbox"/>	

2 TRAUMATIC: CONTACT/COLLISIONS HELMETED SPORTS

	YES	NO	COMMENTS
All contact/collision, helmeted practices and competitions adhere to existing ethical standards.	<input type="checkbox"/>	<input type="checkbox"/>	
All contact/collision, helmeted practices and competitions adhere to keeping the head out of blocking and tackling.	<input type="checkbox"/>	<input type="checkbox"/>	
All contact/collision, helmeted practices and competitions adhere to prohibiting the use of the helmet as a weapon.	<input type="checkbox"/>	<input type="checkbox"/>	
All contact/collision, helmeted practices and competitions adhere to not deliberately inflicting injury on another player.	<input type="checkbox"/>	<input type="checkbox"/>	
All contact/collision, helmeted practices and competitions adhere to maintaining and certifying helmets to existing helmet safety standards.	<input type="checkbox"/>	<input type="checkbox"/>	

3 NON-TRAUMATIC: GENERAL

	YES	NO	COMMENTS
All practices and strength and conditioning sessions adhere to established scientific principles of acclimatization and conditioning.	<input type="checkbox"/>	<input type="checkbox"/>	
Conditioning periods are phased in gradually and progressively to encourage proper exercise acclimatization and to minimize the risk of adverse effects on health.	<input type="checkbox"/>	<input type="checkbox"/>	
The first seven days of any new conditioning cycle are considered a transition period and a time of physiologic vulnerability for athletes.	<input type="checkbox"/>	<input type="checkbox"/>	
Transition periods for athletes include, but are not limited to, returning after an injury or illness.	<input type="checkbox"/>	<input type="checkbox"/>	
Transition periods for athletes include, but are not limited to, returning after school break (e.g., winter, spring, summer).	<input type="checkbox"/>	<input type="checkbox"/>	

NON-TRAUMATIC: GENERAL CONTINUED	YES	NO	COMMENTS
Transition periods for athletes include, but are not limited to, beginning as a delayed start.	<input type="checkbox"/>	<input type="checkbox"/>	
Training and conditioning sessions are appropriately calibrated and include limitations on total volume and intensity of activity, especially during the first four days of transition periods.	<input type="checkbox"/>	<input type="checkbox"/>	
All workouts have a written plan that is exercise science-based, physiologically sport-specific, and tailored to the individual.	<input type="checkbox"/>	<input type="checkbox"/>	
Workout plans are approved by a credentialed strength and conditioning professional, or the responsible sport coach if a strength and conditioning professional is not available at the institution.	<input type="checkbox"/>	<input type="checkbox"/>	
Components of the workout plan include volume, intensity, mode and duration.	<input type="checkbox"/>	<input type="checkbox"/>	
The activity location is stated in the workout plan to accommodate venue-specific emergency action planning.	<input type="checkbox"/>	<input type="checkbox"/>	
Workout plans are reproducible upon request and shared with the primary athletics health care providers (team physician and athletic trainer) before the session in which they are to be used.	<input type="checkbox"/>	<input type="checkbox"/>	
Modification due to hazardous environmental conditions, scheduling considerations, etc., is supported. The amended workout plan maintains the above principles.	<input type="checkbox"/>	<input type="checkbox"/>	
Exercise never is used for punitive purposes.	<input type="checkbox"/>	<input type="checkbox"/>	
Educational background, sport experience and credentialing are verified for all strength and conditioning professionals.	<input type="checkbox"/>	<input type="checkbox"/>	
All strength and conditioning professionals have a reporting line into the sports medicine or sport performance lines of the institution.	<input type="checkbox"/>	<input type="checkbox"/>	
Emergency action plans are developed and rehearsed annually for all venues in which practices or competitions are conducted.	<input type="checkbox"/>	<input type="checkbox"/>	
Emergency action plans are developed and rehearsed annually for head and neck injuries.	<input type="checkbox"/>	<input type="checkbox"/>	
Emergency action plans are developed and rehearsed annually for cardiac arrest.	<input type="checkbox"/>	<input type="checkbox"/>	
Emergency action plans are developed and rehearsed annually for exertional heat illness and heat stroke.	<input type="checkbox"/>	<input type="checkbox"/>	
Emergency action plans are developed and rehearsed annually for exertional rhabdomyolysis.	<input type="checkbox"/>	<input type="checkbox"/>	

NON-TRAUMATIC: GENERAL CONTINUED	YES	NO	COMMENTS
Emergency action plans are developed and rehearsed annually for exertional collapse associated with sickle cell trait.	<input type="checkbox"/>	<input type="checkbox"/>	
Emergency action plans are developed and rehearsed annually for any exertional or non-exertional collapse.	<input type="checkbox"/>	<input type="checkbox"/>	
Emergency action plans are developed and rehearsed annually for asthma.	<input type="checkbox"/>	<input type="checkbox"/>	
Emergency action plans are developed and rehearsed annually for diabetic emergency.	<input type="checkbox"/>	<input type="checkbox"/>	
Strength and conditioning venues have emergency action plans specific to the venue, sport and circumstances.	<input type="checkbox"/>	<input type="checkbox"/>	
The institution has adopted requirements for the annual education and training for the prevention of sudden death in sport for strength and conditioning professionals.	<input type="checkbox"/>	<input type="checkbox"/>	
The institution has adopted requirements for the annual education and training for the prevention of sudden death in sport for sport coaches.	<input type="checkbox"/>	<input type="checkbox"/>	
The institution has adopted requirements for the annual education and training for the prevention of sudden death in sport for athletic trainers.	<input type="checkbox"/>	<input type="checkbox"/>	
The institution has adopted requirements for the annual education and training for the prevention of sudden death in sport for team physicians.	<input type="checkbox"/>	<input type="checkbox"/>	
The institution has adopted requirements for the annual education and training for the prevention of sudden death in sport for collegiate athletes.	<input type="checkbox"/>	<input type="checkbox"/>	
The institution has adopted requirements for the annual education and training for the prevention of sudden death in sport for athletics administrators.	<input type="checkbox"/>	<input type="checkbox"/>	



REFERENCES

1. Yau R, Kucera KL, Thomas LC, Price HM, Wolff TLC, Cantu RC. *Catastrophic sports injury research: Thirty-fifth annual report Fall 1982-Spring 2017*. Chapel Hill, NC: National Center for Catastrophic Sport Injury Research at the University of North Carolina at Chapel Hill; September 25, 2018.
2. Kucera KL, Fortington LV, Wolff CS, Marshall SW, Finch CF. Estimating the international burden of sport-related death: a review of data sources. *Inj Prev*. 2018.
3. Van SC, Bloor CM, Mueller FO, Cantu RC, Olson HG. Nontraumatic sports death in high school and college athletes. *Med Sci Sports Exerc*. 1995;27(5):641-647.
4. Kucera KL, Klossner D, Colgate B, Cantu RC. *Annual survey of football injury research: 1931-2017*. Chapel Hill, NC: National Center for Catastrophic Sport Injury Research at the University of North Carolina at Chapel Hill for the American Football Coaches Association, National Collegiate Athletic Association, & National Federal of State High School Associations.; February 16 2018.
5. Boden BP, Breit I, Beachler JA, Williams A, Mueller FO. Fatalities in high school and college football players. *Am J Sports Med*. 2013;41(5):1108-1116.
6. Cantu RC, Mueller FO. Brain Injury-related Fatalities in American Football, 1945–1999. *Neurosurgery*. 2003;52(4):846-853.
7. Harmon KG, Drezner JA, Klossner D, Asif IM. Sick cell trait associated with a RR of death of 37 times in National Collegiate Athletic Association football athletes: a database with 2 million athlete-years as the denominator. *Br J Sports Med*. 2012;46(5):325-330.
8. Harris KM, Haas TS, Eichner ER, Maron BJ. Sick cell trait associated with sudden death in competitive athletes. *Am J Cardiol*. 2012;110(8):1185-1188.
9. National Athletic Trainers' Association. Consensus statement: Sick cell trait and the athlete. <https://www.nata.org/sites/default/files/SickleCellTraitAndTheAthlete.pdf>. Accessed April 11, 2019.
10. Sick cell solubility test, Division I NCAA Bylaw, §17.1.5.1 (2010).
11. Sick cell solubility test, Division II NCAA Bylaw, §17.1.5.1 (2012).
12. Confirmation of Sick Cell Trait Status, Division III NCAA Bylaw, §17.1.6.4.1 (2013).
13. Adams WM, Casa DJ, Drezner JA. Sport safety policy changes: Saving lives and protecting athletes. *J Athl Train*. 2016;51(4):358-360.
14. Eichner ER. "A Stitch in Time" and "If 6 was 9": Preventing exertional sickling deaths and probing team rhabdomyolysis outbreaks. *Curr Sports Med Rep* 2016;15(3):122-123.
15. Casa DJ, Anderson SA, Baker L, et al. The inter-association task force for preventing sudden death in collegiate conditioning sessions: Best practices recommendations. *J Athl Train*. 2012;47(4):477-480.
16. McGrew CA. NCAA football and conditioning drills. *Curr Sports Med Rep*. 2010;9(4):185-186.
17. Five-day acclimatization period, Division I; Division II NCAA Bylaw, §17.10.2.4 (2003).
18. Five-day acclimatization period, Division III NCAA Bylaw, §17.10.2.2 (2004).
19. Strength and Conditioning Coach first aid / CPR certification and authority of sports medicine staff (FBS / FCS), Division I NCAA Bylaw, §13.11.3.7.4 (2003).
20. Health and safety oversight, Division II NCAA Bylaw, §17.01.2 (2004).
21. Strength and conditioning activities as follows, Division III NCAA Bylaw, §17.02.1.1.1(j) (2011).
22. Strength and Conditioning Coach certification, Division II NCAA Bylaw, §11.1.7 (2016).
23. Strength and Conditioning Coach certification, Division I NCAA Bylaw, §11.1.5 (2014).
24. Mueller FO. Catastrophic head injuries in high school and collegiate sports. *J Athl Train*. 2001;36(3):312-315.
25. Torg JS, Sennett B, Vegso JJ, Pavlov H. Axial loading injuries to the middle cervical spine segment. *Am J Sports Med*. 1991;19(1):6-20.
26. Torg JS, Vegso JJ, O'Neill MJ, Sennett B. The epidemiologic, pathologic, biomechanical, and cinematographic analysis of football-induced cervical spine trauma. *Am J Sports Med*. 1990;18(1):50-57.

27. Anderson S. NCAA Football Off-Season Training: Unanswered Prayers ... A Prayer Answered. *J Athl Train.* 2017;52(2):145-148.
28. Anderson SA. The junction boys syndrome. *J Strength Cond Res.* 2012;26(5):1179-1180.
29. Demartini JK, Casa DJ. Who is responsible for preventable deaths during athletic conditioning sessions? *J Strength Cond Res.* 2011;25(7):1781.
30. The principle of sportsmanship and ethical conduct, NCAA Constitution, §2.4 (1996).
31. Andres K. *2018 and 2019 NCAA Men's and Women's Soccer Rules.* Indianapolis, IN: National Collegiate Athletic Association; 2018.
32. Hyland A. *2018-19 NCAA Men's Basketball Rules.* Indianapolis, IN: National Collegiate Athletic Association; 2018.
33. Redding R. *2017 NCAA Football Rules and Interpretations.* Indianapolis, IN: National Collegiate Athletic Association; 2018.
34. National Operating Committee on Standards for Athletic Equipment. Standard performance specification for newly manufactured football helmets. In. Kansas City, KS: NOCSAE; 2015.
35. National Operating Committee on Standards for Athletic Equipment. Standard performance specification for recertified football helmets. In. Kansas City, KS: NOCSAE; 2015.
36. ASTM International. *Sports equipment, playing surfaces and facilities; pedestrian/walkway safety and footwear; amusement rides and devices; snow skiing.* Vol 15.07. West Conshohocken, PA: ASTM International; 2016.
37. Levy ML, Ozgur BM, Berry C, Aryan HE, Apuzzo ML. Analysis and evolution of head injury in football. *Neurosurgery.* 2004;55(3):649-655.
38. Casa DJ, Guskiewicz KM, Anderson SA, et al. National Athletic Trainers' Association position statement: Preventing sudden death in sports. *J Athl Train.* 2012;47(1):96-118.
39. Garrett AT, Rehrer NJ, Patterson MJ. Induction and decay of short-term heat acclimation in moderately and highly trained athletes. *Sports Med.* 2011;41(9):757-771.
40. Cooper ER, Ferrara MS, Casa DJ, et al. Exertional heat illness in American football players: When is the risk greatest? *J Athl Train.* 2016.
41. Casa DJ, Demartini JK, Bergeron MF, et al. National Athletic Trainers' Association position statement: Exertional heat illness. *J Athl Train.* 2015;50(9):986-1000.
42. Eichner ER. Football team rhabdomyolysis: The pain beats the gain and the coach is to blame. *Curr Sports Med Rep.* 2018;17(5):142-143.
43. Eichner ER. Exertional rhabdomyolysis stays in the news. *Curr Sports Med Rep.* 2016;15(6):378-379.
44. Drezner JA, Chun JS, Harmon KG, Derminer L. Survival trends in the United States following exercise-related sudden cardiac arrest in the youth: 2000-2006. *Heart Rhythm.* 2008;5(6):794-799.
45. Drezner JA, Courson RA, Roberts WO, Mosesso VN, Link MS, Maron BJ. Inter-association task force recommendations on emergency preparedness and management of sudden cardiac arrest in high school and college athletic programs: A consensus statement. *J Athl Train.* 2007;42(1):143-158.
46. Herring SA, Kibler WB, Putukian M. Sideline preparedness for the team physician: A consensus statement-2012 update. *Med Sci Sports Exerc.* 2012;44(12):2442-2445.
47. Concussion safety protocol management. <http://www.ncaa.org/concussionsafety>. Accessed April 11, 2019.
48. IOM (Institute of Medicine). *Strategies to improve cardiac arrest survival: A time to act.* Washington D.C.: Institute of Medicine;2015.
49. Hainline B, Drezner JA, Baggish A, et al. Interassociation Consensus Statement on Cardiovascular Care of College Student-Athletes. *J Am Coll Cardiol.* 2016;67(25):2981-2995.
50. Casa DJ, Armstrong LE, Kenny GP, O'Connor FG, Huggins RA. Exertional heat stroke: new concepts regarding cause and care. *Curr Sports Med Rep.* 2012;11(3):115-123.
51. Parsons JT, ed *2014-2015 NCAA Sports Medicine Handbook.* Indianapolis, IN: National Collegiate Athletic Association; 2014.

APPENDIX A

2016 SAFETY IN COLLEGE FOOTBALL SUMMIT AGENDA

AGENDA

National Collegiate Athletic Association
Safety in College Football Summit

Orlando, Florida

February 10-11, 2016

DAY 1

1. **Welcome and summit overview. (Scott Anderson and Brian Hainline)**
2. **Topic 1: Sensor and clinical data regarding football practice and head exposure.**
 - a. Campus research. (Stefan Duma, Thomas Druzgal, Jacob Marucci, Jason Mihalik)
 - b. Big 12 research. (Scott Anderson, Allen Hardin)
 - c. Roundtable discussion and report out.
 - d. Referendum: Year-round football practice contact.
3. **Topic 2: Catastrophic injury in football.**
 - a. Traumatic. (Kevin Guskiewicz)
 - b. Nontraumatic. (Scott Anderson, Doug Casa)
 - c. Roundtable discussion and report out.
 - d. Referendum: Action plan for mitigating catastrophic injury in football.
4. **Topic 3: Diagnosis and management of sport-related concussion guidelines.**
 - a. Guidelines overview. (Brian Hainline, Scott Anderson)
 - b. Concussion diagnosis and management update: New data from Concussion Assessment, Research and Education Consortium. (Steven Broglio, Thomas McAllister, Michael McCrea)
 - c. Re-examining concussion treatment: Agreements from the TEAM meeting? (Anthony Kontos)
 - d. Roundtable discussion and report out.
 - e. Referendum: Diagnosis and management of sport-related concussion.

DAY 2

1. **Opening remarks. (Scott Anderson and Brian Hainline)**
2. **Topic 4: Independent medical care. (Scott Anderson and Brian Hainline)**
 - a. Roundtable discussion and report out.
 - b. Referendum: Independent medical care.
3. **Topic 5: Interassociation consensus statements.**
 - a. Year-round football practice contact.
 - b. Catastrophic injury in football.
 - c. Diagnosis and management of sport-related concussion.
 - d. Independent medical care.

4. **Closing remarks.**

APPENDIX B

2016 SAFETY IN COLLEGE FOOTBALL SUMMIT PARTICIPANTS

Jeff Allen, Head Athletic Trainer, University of Alabama
(attending on behalf of Nick Saban)

Scott Anderson, College Athletic Trainers' Society,
University of Oklahoma

Doug Aukerman, Pac-12 Conference

Julian Bailes, M.D., Congress of Neurological Surgeons,
American Association of Neurological Surgeons

Stevie Baker-Watson, Director of Athletics,
DePauw University

Brad Bankston, Commissioner, Old Dominion
Athletic Conference

Karl Benson, Commissioner, Sun Belt Conference

Bob Boerigter, Commissioner, Mid-America
Intercollegiate Athletics Association

Bob Bowlsby, Commissioner, Big 12 Conference;
Chair, Football Oversight Committee

Matthew Breiding, Centers for Disease Control
and Prevention

Steve Broglio, M.D., Principal Investigator,
CARE Consortium, University of Michigan

William Bynum, President, Mississippi Valley State
University

Jeff Bytomski, D.O., American Osteopathic Academy
of Sports Medicine

Carolyn Campbell-McGovern, The Ivy League

Doug Casa, Ph.D., Consortium Director, Division on
Exertional Injury, National Center for Catastrophic Sport
Injury Research; Chief Executive Officer, Korey Stringer
Institute; Director, Athletic Training Education,
University of Connecticut

Bob Casmus, Committee on Competitive Safeguards and
Medical Aspects of Sports, Catawba College

Scott Caulfield, National Strength and Conditioning
Association

Randy Cohen, National Athletic Trainers' Association

Bob Colgate, National Federation of State High School
Associations

Dawn Comstock, Associate Professor, University of
Colorado, Denver

Kevin Crutchfield, M.D., American Academy of Neurology

Ty Dennis, Division II Student-Athlete Advisory
Committee, Minnesota State University, Mankato

Jon Divine, M.D., President, American Medical Society
for Sports Medicine

Tom Dompier, Ph.D., President, Datalys Center for
Sports Injury Research and Prevention

Jason Druzgal, M.D., Neuroradiologist, University
of Virginia

Stefan Duma, Ph.D., Director, School of Biomedical
Engineering and Sciences, Virginia Polytechnic Institute
and State University

Ruben Echemendia, Ph.D., President, Sports
Neuropsychology Society

Brent Feland, M.D., Collegiate Strength and Conditioning
Coaches Association

Scott Gines, Director of Athletics, Texas A&M University-
Kingsville

Kevin Guskiewicz, Ph.D., University of North Carolina,
Chapel Hill

Allen Hardin, Senior Associate Athletics Director,
University of Texas at Austin

Steven Hatchell, President, National Football Foundation

Bill Heinz, Chair, Sports Medicine Advisory Committee,
National Federation of State High School Associations

Jaime Hixson, Associate Commissioner, Mountain West
Conference

Peter Indelicato, American Orthopaedic Society for
Sports Medicine

Nick Inzerello, Senior Director, Football Development,
USA Football

Jay Jacobs, Division I Strategic Vision and Planning
Committee, Auburn University

Chris Jones, Division I Football Oversight Committee
(proxy), University of Richmond

Kerry Kenny, Assistant Commissioner,
Big Ten Conference

Zachary Kerr, Director, Datalys Center for Sports
Injury Research and Prevention

Anthony Kontos, Ph.D., Assistant Research Director,
Sports Medicine Concussion Program, University
of Pittsburgh Medical Center

William Lawler, Southeastern Conference

Josephine Lee, Board Member, College Athletic
Trainers' Society

Donald Lowe, Board Member, College Athletic
Trainers' Society

Jack Marucci, Louisiana State University

Thomas McAllister, M.D., Principal Investigator,
CARE Consortium

Michael McCrea, Ph.D., Principal Investigator,
CARE Consortium

William Meehan, M.D., American Academy of Pediatrics

Jason Mihalik, Ph.D., University of North Carolina,
Chapel Hill

Bob Murphy, Board Member, College Athletic
Trainers' Society

Bob Nielson, Chair, NCAA Football Rules Committee
Scott Oliaro, Board Member, College Athletic Trainers' Society
Kene Orjioke, Division I Student-Athlete Advisory Committee, University of California, Los Angeles
Steve Pachman, J.D., Montgomery McCracken
Julie Cromer Peoples, Senior Woman Administrator, University of Arkansas, Fayetteville
Sourav Poddar, M.D., American College of Sports Medicine
Kayla Porter, Division III Student-Athlete Advisory Committee, Frostburg State University
Rogers Redding, Secretary-Rules Editor, NCAA Football Rules Committee
Yvette Rooks, Board Member, College Athletic Trainers' Society
Eric Rozen, Board Member, College Athletic Trainers' Society
Scott Sailor, President, National Athletic Trainers' Association

Jon Steinbrecher, Commissioner, Mid-American Conference
Ken Stephens, National Operating Committee on Standards for Athletic Equipment
Edward Stewart, Senior Associate Commissioner, Big 12 Conference
Michael Strickland, Senior Associate Commissioner, Atlantic Coast Conference
Grant Teaff, Executive Director, American Football Coaches Association
Buddy Teevens, Coach, Dartmouth College
James Tucker, M.D., Board Member, College Athletic Trainers' Society
Steve Walz, Associate Director of Athletics, University of South Florida
Alfred White, Senior Associate Commissioner, Conference USA

STAFF PARTICIPANTS

Brian Burnsed, Associate Director, Communications
Dawn Buth, Associate Director, Sport Science Institute
Cassie Folck, Coordinator, Sport Science Institute
Brian Hainline, Chief Medical Officer, NCAA
Kathleen McNeely, Chief Financial Officer, NCAA
Terrie Meyer, Executive Assistant, Sport Science Institute

John Parsons, Director, Sport Science Institute
Chris Radford, Associate Director, Public and Media Relations
Stephanie Quigg, Director, Academic and Membership Affairs

APPENDIX C

ENDORISING ORGANIZATIONS

The following organizations have endorsed this document:

- American Association of Neurological Surgeons
- American Medical Society for Sports Medicine
- American Orthopaedic Society for Sports Medicine
- American Osteopathic Academy of Sports Medicine
- College Athletic Trainers' Society
- Collegiate Strength and Conditioning Coaches Association
- Congress of Neurological Surgeons
- Korey Stringer Institute
- National Athletic Trainers' Association
- National Strength and Conditioning Association
- National Operating Committee for Standards on Athletic Equipment
- Sports Neuropsychology Society

The following organization has affirmed the value of this document:

- American Academy of Neurology





SPORT SCIENCE
INSTITUTE™

PREVENTING CATASTROPHIC INJURY
AND DEATH IN COLLEGIATE ATHLETES | JULY 2019

